

Comparing the impact of chronic energy restriction and vitamin E supplementation on the behavior of adult rats

Derlange B. Diniz, Suzana L. de Oliveira, Liana L. Melo, Jaime Amaya-Farfan

Food Security and Nutrition Laboratory, State University of Ceará, Fortaleza, CE, Brazil, Faculty of Nutrition, Federal University of Alagoas, Maceió, AL, Brazil, University of São Francisco/Bragança Paulista, SP, Brazil, Department of Food and Nutrition, University of Campinas - Unicamp, Brazil

SUMMARY. The purpose of this work was to investigate the influence of energy restriction and vitamin E supplementation on memory, learning, anxiety and spontaneous locomotion in adult rats. Three-month old male Wistar rats were grouped according to diet: Control (AIN 93-M; n=18), Supplemented (AIN 93-M + 1425 IU all-*rac*- α -tocopheryl acetate/kg diet; n=22) and Restricted (AIN 93-M with 30% reduction in carbohydrate energy; n=23). Sixteen weeks after, the passive avoidance (PA), elevated plus-maze (EPM) and open field (OF) tests were applied. In the EPM test, the behavioral profile of the supplemented group was characterized by a lower frequency of entries into the open arms ($P < 0,026$), whereas the restricted group showed a lower frequency of head dipping ($P < 0,003$). The ratio between the time span of the shocks and the number of attempts were larger for the supplemented than for the non-supplemented animals ($P = 0,0474$), thus suggesting a delay in learning in the PA test. Taken together, these results suggest that a long-term combination of carbohydrate energy restriction in rats should not cause negative behavioral alterations. Compared with vitamin E supplementation, the restricted diet performed equally or better in rats as an alternate antioxidant diet.

Key words: Diet composition, behavior, energy restriction, vitamin E supplementation, antioxidant diet.

RESUMEN. Comparación del impacto entre la restricción energética crónica y la suplementación con vitamina E en el comportamiento de ratas adultas. Se investigó la influencia de la restricción energética en comparación a la suplementación con vitamina E en la memoria, aprendizaje, ansiedad y locomoción espontánea en ratas adultas. Machos Wistar de tres meses de edad fueron agrupados de acuerdo a las dietas: Control (AIN 93-M, n=18), Suplementados (AIN 93-M + 1425 UI all-*rac*- α -tocoferil acetato / kg de dieta; n=22) y Restrictos (AIN 93-M con 30% de restricción energética en los carbohidratos; n=23). Dieciséis semanas después, fueron aplicadas las pruebas de esquivia pasiva (PA), laberinto elevado en cruz (EPM) y de campo abierto (OF). En la prueba de EPM, el grupo suplementado mostró una menor frecuencia de entradas en los corredores abiertos ($P < 0,026$), mientras el grupo restricto registró menor frecuencia de bajadas de cabeza ($P = 0,003$). La razón entre la duración de los choques y el número de intentos fue superior para el grupo suplementado, que para los no suplementados ($P = 0,0474$), lo cual sugiere un leve perjuicio en el aprendizaje de los suplementados, según la prueba de PA. Tomados en conjunto, estos resultados sugieren que la restricción crónica en energía de carbohidratos no provoca alteraciones negativas en el comportamiento de ratas adultas y que sus beneficios pueden superar los obtenidos al suplementar la dieta patrón con vitamina E, como opción de dieta antioxidante.

Palabras clave: Dieta, comportamiento, restricción energética, suplementación con vitamina E, dieta antioxidante.

INTRODUCTION

The impact of the modern environment on well-being, including the diet, is a main concern of health authorities throughout the world. Special emphasis has been placed on the use of antioxidants, whether as part of the diet or in the form of supplements in order to minimize the impact of environmental factors on the general state of health. While no official reference intakes on the vast list of antioxidants have been issued, and as more antioxidants, both synthetic and natural, become available in the market, conscious consumers may prefer to adopt one of several forms of food restriction, among which carbohydrate energy restriction has met some success in the past, as an option to provide themselves with a share of antioxidant benefits.

Energy restriction has been recognized as one of the most consistent dietary manipulations to delay aging in mammals (1-3). Some researchers have focused on the benefits of this practice on the physical performance, learning and information retrieval capacity of old rats (4-6), the learning capacity of young rats (7) and on behavioral tests using experimental models for degenerative diseases (8,9). On the other hand, potentially harmful effects of energy restriction on brain development, such as the effect of protein-energy deficits on cognitive and motor performance in young animals, have also been pointed out (10-13). Although the beneficial effects attributed to energy restriction depend on the adequacy level of all the essential nutrients supplied by the diet and, therefore, defining the diet composition may be of fundamental

importance to the experimental design (7), this issue has been poorly addressed.

The health properties of vitamin E have been extensively recognized, including inhibition of smooth-muscle cell proliferation, preservation of endothelial function, inhibition of monocyte-endothelial cell adhesion, inhibition of monocyte reactive oxygen species and cytokine release, and inhibition of platelet adhesion and aggregation. These cellular responses to α -tocopherol are associated with transcriptional and post-transcriptional events. Activation of diacylglycerol kinase and protein phosphatase 2A, and the inhibition of protein kinase C, cyclooxygenase, lipooxygenase, tyrosine kinase 2-phosphorylation, and cytokine release by α -tocopherol are all examples of post-transcriptional regulation that explain the vitamin's antioxidant actions in the body (14), while its consumption as an antioxidant supplement gains acceptance owing to its very low toxicity (15).

The brain contains high levels of unsaturated lipids and is responsible for 20 to 25% of the body's oxygen consumption, under conditions that favor the occurrence of oxidative stress. Quantitatively, vitamin E is the predominant encephalic lipophilic antioxidant, whose main function is to protect brain lipids from oxidative damage (16). The physiological decline associated with aging, including psychomotor and cognitive disorders, is considered to be partly the result of an accumulation of molecular oxidative damage (17,18), and this process may be ameliorated by antioxidant therapies, over and above the general recommendation of a plethora of antioxidant foods.

Because the brain tissue is particularly susceptible to oxidative stress we have compared a diet protocol of carbohydrate energy restriction with vitamin E supplementation in Wistar rats and found that both can offer similar protection against oxidative damage to various tissues, except to the brain, where the restricted diet had a significantly better performance (19).

Despite the potential benefits attributed to the above mentioned diet models, the possible repercussions of such interventions for prolonged periods of time on adult animal behavior have not been thoroughly studied and should merit a comparative evaluation. Therefore, the objective of the present study was to examine the nutritional-behavioral relationship, specifically regarding the anxiety level, locomotion activity, learning and memory of adult rats submitted to chronic consumption of energy restricted, as compared to vitamin-E supplemented diets. For this purpose, the classical open field (OF), elevated-plus maze (EPM) and passive avoidance (PA) tests were used.

MATERIALS AND METHODS

Animals and diets

Sixty-three 21-day old male Wistar rats (Vivarium Center, University of Campinas, Brazil) were housed in collective

cages ($22 \pm 2^\circ\text{C}$; 12h light-dark cycles) and fed on commercial laboratory chow (Labina, Ralston-Purina do Brasil, Ltd., Campinas, Brazil) *ad libitum* for eight weeks. At three months of age, the animals were transferred to individual cages to start the experiment and randomly allocated to one of three diet groups. Group *C* (control; $n=18$) had free access to the maintenance AIN 93-M diet (20), while group *S* (supplemented; $n=22$) was fed diet *C ad libitum*, but with the addition of 1425 IU of vitamin E (all-*rac*- α -tocopherol acetate), and group *R* (restricted; $n=23$) received a modified *C* diet at a lower amounts and with less carbohydrate, as detailed in Table 1. Initially, all groups underwent a one-week adaptation period consuming the control diet.

TABLE 1
Composition of the three diets based on the recommendations of AIN 93-M*

Ingredients	Diets (g/100g)		
	Control	Restricted	Supplemented
Corn starch (87,6%)	62.07	54.15	62.07
Sucrose (99,5%)	10.00	7.22	10.00
Soybean oil	4.00	5.71	4.00
Fiber (cellulose)	5.00	5.86	5.00
AIN-93 M mineral mixture [†]	3.50	5.00	3.50
AIN-93 M vitamin mixture [‡]	1.00	1.43	1.00
L-cystine	0.18	0.26	0.18
Choline bitartrate	0.25	0.36	0.25
<i>Tert</i> -butylhydroquinone	0.0008	0.0011	0.0008
all- <i>rac</i> - α -tocopherol acetate (0.5 α -TE ^{**})	—	—	0.285 ^{**}
Total energy (joules)	1454.82	1454.44	1454.82

*Reeves et al. (20). [†]contains 20.98g sucrose/100g.

[‡]contains 97.47g sucrose/100g.

** α -TE:equivalents of α -tocopherol; 1 α -TE=1 mg *d*- α -tocopherol.

^{**}sufficient to supply additional 1425IU/kg of diet (or a total of 1500 U/kg).

The restricted diet was formulated such that when offering 70% of the amount consumed by group *C*, the amounts delivered for all nutrients would be the same as for group *C*, with the exception of the carbohydrates. Consequently, by restricting diet consumption (or energy intake) to 70% of that of the controls, the *R* group was ingesting 40% less carbohydrate. Diet *S*, on the other hand, supplied twenty times the amount of vitamin E recommended for the rat (20). In order to keep a closer check on energy consumption, the total energy of the diets was determined using a Parr calorimeter, model 1261/1563 (Moline IL, USA).

The feeding trial was carried out for 16 weeks and the entire study was performed according to the Guide to the Care and Use of Experimental Animals of the University of Campinas, Brazil. The amount of diet to be fed to the *R* group was determined from the daily average consumed by the *C* group. Body weight changes were monitored and recorded weekly.

Behavioral tests

At the end of the 16th week, the effects of the diet on spontaneous motor activity, anxiety level, learning capacity and memory were measured by applying the open field (OF), elevated plus-maze (EPM) and passive avoidance (PA) tests, respectively. The tests were conducted in that sequence throughout a period of two weeks, with intervals of three days between tests. Preceding every test, the freshly fed animals were allowed to stay in the testing environment for 10 minutes in order to remove the effects of the natural search for food or of inhibition due to a new environment. All procedures were manually and video recorded by the researcher from a location visually inaccessible to the subjects.

For the OF test, the instrument consisted of a transparent acrylic chamber (100x100x40cm) with 25 square plots drawn on the floor. The animal was initially placed in one of the corners and, during a 5-min observation period, peripheral and central crossings and rearing, self-washing and defecation activities were registered.

The EPM test was conducted in a roofless, Formica-coated wooden device, positioned 50cm above the laboratory floor and consisting of two open arms (50x10x1cm), with acrylic side walls (1cm), and two closed (50x10x40cm) arms. In order to start the test, the animals were placed at the center of the maze, facing one of the closed arms. Their behavior was observed and recorded for 5min. Standing times in the open and closed arms, self-washing routines, protected (head projected towards the closed arms or central platform) and unprotected (head projected towards open arms), head dipping, listening and both protected and unprotected cautious exploration attempts, were recorded.

The device for the PA test was a box (40x22x30cm) with a metal roof and walls and a floor with stainless steel bars (0.3 cm in diameter, spaced 1.2 cm from each other) connected to a low intensity (1mA) shock generator, to provoke an adverse reaction. A Formica platform (8 x 13 cm), hoisted on stilts, was placed inside the box to one side. The front of the box was of transparent acrylic material to allow for the researcher to see the animal and make a video recording. The test consisted of two phases: acquisition and retention. In the first phase the animal should remain on the platform where he was placed at the start of the test, avoiding the shocks for 200 sec, as a learning parameter. The number of descents (attempts), the time taken to descend from the platform (latency), and the time remaining on the metal bars (receiving shocks) were recorded. Animals remaining on the bars receiving shocks for more than 300 sec were eliminated from the experiment. The second phase, carried out 48 h later, tested retention of the acquired behavior, measuring the time required for the animal to descend from the platform and place his four paws on the bars, without shocks (maximum waiting period, 600 sec). In this phase the number of false descents from the platform was also measured (placing

the front paws on the bars and returning to the platform) and the number of rearing movements.

Alpha-Tocopherol in brain tissue

This vitamin was determined by high performance liquid chromatography (HPLC) according to Sharma & Kumar (21), with modifications. The brain tissue was homogenized (50 g/L; 0.05M phosphate buffer, pH 7.4; 90 μ M butylhydroxytoluene -BHT) in an ice bucket, using a homogenizer with a Teflon pestle. 400 μ L of absolute ethanol were added to the brain homogenate and the mixture vortexed for 1 minute. After centrifugation (12,000 x g for 5 minutes), the tocopherol was recovered in methanol following extraction with *n*-hexane. Elution from the HPLC equipment (Varian, model 9012, equipped with a 9075 fluorescence detector) was monitored at 290 (excitation) and 330nm (emission). A standard calibration curve for (\pm)- α -tocopherol was used for quantification, and recorded per unit protein in the tissue. With a few exceptions, the alpha-tocopherol determinations were carried out in half of the animals of each group.

Statistical analyses

The characteristics of normality of distribution of the data and homogeneity of the variances were analyzed using the tests of Lilliefors and Levene, respectively. Whenever applicable, monovariate ANOVA was performed and differences between groups detected by the tests of Snedecor and Tukey-HSD. When the criteria for parametric analysis were not met, the differences were evaluated by the Kruskal-Wallis test, adopting the Nemenyi test for multiple comparisons. A level of $p \leq 0.05$ was selected to indicate minimum significance for all comparisons. The Statistica for Windows program (StatSoft Inc. Tulsa, OK/USA) was used for the statistical analyses.

RESULTS

Body weight and brain α -tocopherol levels

As expected, the mean weight gain of the restricted group at the end of the experimental period was significantly lower, or approximately 22% of the weight gained by the others (initial mean weights: C = 323.4 \pm 25; S = 329.8 \pm 28; R = 332.0 \pm 26g; final mean weights: C = 467.1 \pm 40; S = 473.6 \pm 52; R = 364.7 \pm 17g; maximal $P < 0.001$ for changes). The respective final mean brain α -tocopherol concentrations for the control, restricted and supplemented groups were 0.32 \pm 0.02, 0.28 \pm 0.03 and 0.47 \pm 0.05 μ g/ mg protein (P between S and the others < 0.0001).


Open Field

The standard Open Field test is commonly used to assess locomotor, exploratory and anxiety-like behavior in laboratory rats. This test is particularly useful in evaluating the

effects of anxiolytic and anxiogenic drugs, locomotor responses to drug and as well as behavioral responses to novelty.

The results shown in Figure 1 indicate no differences in general locomotion activity (central + peripheral crossings) between the supplemented and restricted animals when compared to the control group, or among themselves ($F(2,58) = 1.35$; $P = 0.2678$), after a five-minute period. Likewise, no alterations were detected in central roaming ($F(2,58) = 1.01$; $P = 0.3694$). Nevertheless, the vitamin-E supplemented group performed a significantly lower number of peripheral crossings during the second minute of the test, with respect to the other two diet groups ($F(2,58) = 5.85$; $P = 0.0048$). As time elapsed, however, differences between groups tended to disappear.

FIGURE 1




Peripheral (I) and Central (II) crossings in the Open Field Test. (Control diet, $n = 17$; Vitamin-E supplemented diet, $n = 22$; Restricted diet, $n = 22$). Asterisk indicates $P < 0.01$ in relation to the control. Values represent the mean and standard error of the mean.

There was also no difference between the groups regarding the total number of rearing attempts ($C: 27.35 \pm 2.41$, $R: 25.2 \pm 1.9$ and $S: 22.1 \pm 2.22$; $F(2,58) = 1.63$; $P = 0.2047$) or self-washings ($C: 0.35 \pm 0.19$, $R: 0.36 \pm 0.28$ and $S: 0.45 \pm 0.21$; $H(2, N = 61) = 2.2414$ $P = 0.3261$), although the number of fecal droppings were significantly greater for the restricted group than for the other two diet groups ($C: 1.18 \pm 0.32$, $R: 3.09 \pm 0.46$ and $S: 0.46 \pm 0.55$; $H(2, N = 61) = 6.54$ $P = 0.0380$).

Elevated-plus maze


This test is used to assess the anxiety produced by any external or internal factor, such as drug, food or environmental perturbation and is based on the animal's preference for dark, closed, rather than bright, open spaces. Statistical analysis of the results of the EPM (Figure 2) indicated that the diets under evaluation produced no differences in the general parameters analyzed by this anxiety test. Nevertheless, excluding any zero-entry events (Figure 3), the analysis showed that animals of the supplemented group entered the open arms a lower number of times, in comparison to the controls ($F(2,38) = 4.04$; $P < 0.026$), suggesting they felt less confident when being in the open. A significantly lower number of head dips ($C: 8.0 \pm 0.98$, $R: 3.29 \pm 0.53$ and $S: 6.64 \pm 1.34$; $F(2,38) = 6.79$; $P < 0.003$), and unprotected dips ($C: 5.42 \pm 0.1$, $R: 1.93 \pm 0.47$ and $S: 3.14 \pm 0.88$; $H(2,0 N = 41) = 7.48$; $P = 0.02$) were also evident for the restricted group, as compared to the control. Head dipping is another indication of self confidence and desire to explore. With respect to listening activities (rearing movements) ($C: 0.94 \pm 0.3$, $R: 0.65 \pm 0.2$ and $S: 0.59 \pm 0.2$) and cautious exploration ($C: 2.06 \pm 0.6$, $R: 1.39 \pm 0.3$ and $S: 2.23 \pm 0.4$), no significant differences were found between the groups under study ($P > 0.05$).

FIGURE 2



Time spent (%) in the Open Arms (I), at Center (II), Number of Entries in the Open (III) and in the Closed (IV) of the animals in diets (C, control, $n = 17$; R, restricted, $n = 23$; S, vitamin-E supplemented, $n = 21$) in the Elevated-Plus Maze test. Values represent means and standard errors of the mean. Equal letters indicate no statistical difference between means ($p > 0.05$).

FIGURE 3




Time spent (%) in an Open Arm (I), at Center (II), Entries in the Open (III) and Entries in the Closed (IV) of animals in diets (C, control, n = 12; R, restricted, n = 15; S, vitamin-E supplemented, n = 14) in the Elevated-Plus Maze test. Animals that did not enter the Open Arms were excluded. Values represent means and standard errors of the mean. Equal letters indicate no statistical difference between means ($p > 0.05$).

Passive avoidance

Passive-avoidance response (PAR) is extensively used for the screening of drugs affecting learning and memory. The test involves training rodents to avoid punishment (normally an electric shock) by curbing a normal behavior (such as the exploratory behavior). At specified intervals after training, the animals are tested again for retention of such learning. The conventional and the most widely employed parameters are step-down latency (SDL) and step-down errors (SDE). In order to test the consistency of the learned behavior, we introduced an extra parameter, namely, the total time spent by the animal in the shock zone (TSZ). The parameters of time-exposure to shock and number of attempts recorded during the avoidance response-acquiring phase (Figure 4) showed no difference between the groups when analyzed individually ($F(2.57) = 0.43$; $P = 0.6529$ and $H(2, N = 60) = 0.88$; $P = 0.6452$, respectively). Since the means did not differ considerably, a new parameter was calculated by taking the ratio between the former two parameters (Figure 4-IV), and this time the ratios showed a significant difference between the supplemented and both control and restricted animals. One rat from the control group and another from the supplemented group were excluded during the first phase of the test for spend-

ing longer than 300 seconds on the bars, without acquiring the avoidance response. In the second phase of the test, no statistical difference was found between the different diet groups with respect to the time spent on the platform ($H(2, N = 61) = 0.88$; $P = 0.6441$), suggesting a similarity in the retention of the learned behavior. Alterations were also not detected with respect to rearing and false descent movements (SDE; Figure 5).

FIGURE 4




Passive Avoidance Response. Means and SEM of duration of shock (min) (I), Number of Attempts (II), Latency (min) to step down from platform (III) and ratio of duration of shock (min)/ attempts (IV) of animals in diets (C, control, n = 17; R, restricted, n = 23; S, vitamin-E supplemented, n = 21) in the Inhibitory Avoidance test. Values represent means and standard errors of the mean (I, II, III) or mean scores (IV). Equal letters indicate no statistical difference between means ($p > 0.05$).

DISCUSSION

The majority of studies on the effect of diet on animal behavior have featured the repercussions of malnutrition and undernutrition on growth and development (11,13,22-25), while others have evaluated the effects of isolated nutritional factors, mainly regarding the benefic action on chronic diseases and aging (2,5,17,18,26-30). Meanwhile, other works have dealt with the influence of energy restriction on development and reproduction (7,8), as well as alternative antioxidant diets (17). This work sought to compare the impact of the chronic administration of two antioxidant dietary regimes, energy restriction (in both carbohydrate content and total energy intake restriction) and vitamin E supplementation on rat behavior.

FIGURE 5



Latency in stepping down from platform – Retention phase (I); Rearings (II) and false stepdowns (III) of animals in diets (C, control, $n = 17$; R, restricted, $n = 23$; S, vitamin-E supplemented, $n = 21$) in the Inhibitory Avoidance test. Values represent means and standard errors of the mean. Equal letters indicate no statistical difference between means ($p > 0.05$).

Upon confrontation of the results obtained in the open field test of the present study with those of the behavioral standard designed by Huston & Bures (31) for adult rats, the observed performance was similar to that described in the literature, with no differences in general activity found between the two diet models investigated. The supplemented animals, however, explored less the peripheral squares between the first and second minute of observation (Figure 1) and this could be somewhat relevant if we were to consider quick responses, but not if we consider the total time interval of the test, when no difference was found among the groups ($C = 72.4 \pm 6$; $R = 74.9 \pm 8$; $S = 60.7 \pm 6$ $F(2,58) = 1.35$; $P = 0.2678$). Since the control and the supplemented groups bore similar body weights and the controls showed a tendency to normal productive locomotion, the possibility that the supplemented animals were discouraged from any greater exploratory activity because of their weight had to be discarded. Therefore, it appears that supplementing the standard diet with extra doses of vitamin E may have resulted in a partial loss of exploratory desire of the animals so treated.

Similarly, McDonald et al. (17) did not observe significant improvement of the psychomotor performance of 24-month old mice fed antioxidant diets containing either vitamin E,

coenzyme Q or a mixture of both for 14 weeks. Interestingly, these authors did notice a moderate impairment in the performance at the initial phase of coordinated running, with either the vitamin alone or in combination with the coenzyme. Sumien et al. (32) in turn have reported deleterious effects in vitamin-E supplemented aged mice, yet not in young ones, when submitted to tests demanding coordination and fast responses, after a short-term supplementation with a dose similar to the one used in this work. Contrasting with our experiment, the supplementation by Sumien et al. (32) and McDonald et al. (17) started when the animals were at least 20 months old.

On the other hand, Shukitt-Hale et al. (30) reported beneficial effects of some antioxidant diets tested by the group, including age-related cognitive behavioral deficits and changes in neuronal signal transduction due to vitamin E supplementation. One of the diets tested was a blueberry based diet that reverted deficits in motor performance. In another study involving the administration of diets with various concentrations of vitamin E and C, plus selenium, fed to 12-month old mice for six months, Richwine et al. (18), also concluded that the diet containing high levels of antioxidants showed to be beneficial, considering that they improved the psychomotor coordination of the 18-month old animals in the rod walking test. The results of our study, therefore, should not necessarily agree with those discussed above for our objective was to assess the response of rats with clearly defined diets and without the intervention of complex mixtures of foods or the cocktail effect of multiple nutrient supplementation. It is still possible, however, that clear-cut beneficial behavioral effects of such antioxidant diets are somewhat age-related, being more evident at the onset of the psychomotor decline, rather than at an early age when target features of behavior can be overridden by either strong adaptive metabolic pathways of the young, or later on, by irreversible metabolic developments that have already been installed.

It is worth mentioning that apart from establishing the spontaneous activity pattern of the animal, the open field test can be used as a simple evaluation of the emotional state. It is known that changes in emotional state are accompanied by various vegetative phenomena, such as changes in the heart rate, voltaic response of the hair and dilatation of the pupils. In addition to these features, one can evaluate the autonomic defecation function, considering that animals that roam less and defecate more are referred to as being more emotionally affected than those with the opposite behavior (31).

Generally and under standard conditions, the behavior of adult rats shows a high negative correlation between defecation and exploration of the central squares. Although the restricted animals excreted a greater number of fecal droppings, it was surprising to find no correlation between this parameter and central roaming. Harrison & Archer (1)

used this test in various assays with mice and observed a greater activity in adult and elderly animals fed a restricted-energy diet, as compared to controls with free access to food. However, more recently, Wu and coworkers reported no detectable differences in the exploratory activity of mice after three (7) and six (8) months consuming diets with 20 and 35% food restriction, respectively. Therefore, it was not surprising to find only a loose correlation between the defecation parameter and the central roaming activity.

The elevated-plus maze is a classical test that measures the degree of anxiety of the animal as being directly proportional to the length of time it spends in the enclosed arms, or avoiding being exposed in the open. With regard to the behavioral features analyzed by the EPM, we observed that, as with the open field results, it was not possible to detect any differences in exploratory or other ethological performance of the adult animal between energy restriction and vitamin E supplementation during the 16-week period, indicating at first glance that neither of the antioxidant dietary models was likely to interfere with the anxiety levels of the animals by this criterion.

It is important to observe, however, that a considerable percentage of the animals, similar for all three groups, did not explore the open arms of the maze at all (C=30%, R= 35%, S=35%), thereby increasing the dispersion of the data in all groups. For this reason the data were further analyzed, this time excluding the non-responsive animals (Figure 3). Upon new analysis, the vitamin-E supplemented group showed a lower percentage of entries in the open arms when compared to the control, thus suggesting an anxiety response in this group, about 28% greater than in the control. The restricted group meanwhile, did not differ from the control. The reason why none of the groups showed a fuller responsiveness is not known. Reactive oxygen species have been critically implicated in various types of pain, including neuropathic and inflammatory pain, in which vitamin E seems to play an analgesic role (33). If this were the case, a slight attitude of indifference could have lowered the response to the test by these animals.

Rodgers & Cole (34), however, examined the behavioral response of mice in the EPM test soon after an experience of aggression, and demonstrated the importance of social stress factors in the production of analgesia. The anxiety response effect of social competition was partially reproduced in mice simply by exposing them to the smell of an aggressive male, producing a reduction in the total number of entries and rearing movements, and also in the percentage of entries into the open arms, as well as the time spent there. By establishing a parallel with the present study, and interpreting the prolonged isolation of the animals as a factor of stress (25), we should arrive at the opposite conclusion because the excess vitamin should confer an additional edge of relaxation to the supplemented group. Therefore, it seems reasonable to suspect that a

factor, other than solely analgesia, may have contributed to the lower number of entries of the supplemented rats in the open arms.

An analysis of the results obtained in some studies evaluating the learning capacity and memory of animals fed diets with different compositions showed that for the inhibitory passive avoidance the behavioral changes more likely to occur in undernourished animals were lower thresholds to the perception of shock, shorter avoidance latency and greater resistance to extinguishing the avoidance response (12,35).

Additionally, Goodrick et al. (36) asserted that once learned, the information remained available for an extensive period of time. These researchers observed a substantial retention of learning up to 45 days after the avoidance test was performed in Wistar rats, both young and old.

In general, undernutrition does not affect acquisition of the avoidance response, but animals in this condition normally require a greater number of attempts on the springboard and in the come-and-go box (12). In our experiment, results of the retention phase (performed 48h later) suggested that neither high levels of vitamin E supplementation or 30% energy restriction for the prolonged period of the study influenced memory recall of the adult rats because in all cases the acquired response was evidently retained.

Moreover, Socci et al. (5), upon administration of a combination of intra-peritoneal injections of phenyl- α -tert-butyl nitron (PBN; 32 mg/kg body weight) and α -tocopherol (200 mg/kg body weight) for approximately 4 to 5 months, in 24 month-old rats, showed that the animals treated with antioxidants had a greater learning acquisition rate. In the experiment of McDonald et al. (17) the authors reported a beneficial effect on learning and memory consolidation of aged rats from feeding a combination of vitamin E and coenzyme Q, but this effect was not detected when the supplemental nutrients were administered individually. In such case, it was hypothesized that a synergic action could have contributed to the final antioxidant effect of the mixture.

In the passive avoidance test, one could think that the vitamin E supplement could have had some influence on the animals' performance because of an induced opioid-type analgesia. This mechanism has been little investigated, but there is evidence to this effect in studies published by Kryzhanovskii et al. (37,38) in the former Soviet Union, where an analgesic action was perceived when vitamin E was administered to women with dysmenorrhea.

The effect of 100 mg α -tocopherol acetate, via intra-muscular injection, on reducing pain, was effective 15-20 minutes after application. When a dose of 500-600 mg was applied during two consecutive menstrual cycles, a residual effect of up to six months after administration was found. The authors showed evidence of a possible involvement of the endogenous opioid system in the analgesic effect induced, mo-

bilizing β -endorphin from the pituitary gland. The involvement of this system was confirmed by injecting naloxone – an antagonist of morphine receptors and endogenous opioid peptides. The analgesic effect of the repeated administration of α -tocopherol has been also blocked after injecting naloxone (37,38). Direct evidence for this association, however, could be sought in future investigations.

In spite of all these considerations, it appears as though any possible handicap associated with the supplemented diet, in relation to energy restriction, may be of such complex nature that the present tests are not designed to determine if the vitamin's analgesic property is involved.

CONCLUSION

By the application of three classical behavioral tests we can conclude that young adult rats submitted to a chronic (four month) carbohydrate-energy restricted diet, suffered no detectable alterations in their normal behavioral characteristics, as evaluated by locomotor activities and various manifestations of anxiety, in comparison to the group fed a standard diet *ad libitum*. The carbohydrate-energy restricted animals performed equally or somewhat better than cohorts that received vitamin E supplementation at a level 20 times the recommended intake. Therefore, it can be said that in addition to performing better than vitamin-E supplementation as a long term antioxidant diet, the prolonged practice of energy restriction, combined with specific carbohydrate limitation, may not affect the normal behavior of the adult rat. Albeit a possible analgesic effect could have played a role in the response of the vitamin-E supplemented group, we think this is an issue that should deserve further investigation.

ACKNOWLEDGMENTS

The authors thank the Brazilian agencies CNPq and CAPES for the doctoral fellowships and to Prof. Dr. Cyro Rêgo Cabral Jr., of the School of Nutrition, Federal University of Alagoas for his kind support with the statistical treatment of the data.

REFERENCES

- Harrison DE, Archer JR. Genotype modulates the effects of caloric restriction on aging process in rodents. Effects of caloric restriction on aging phenomena. In: Ingram DK, Baker GT, Shock NW, eds. The potential for nutrition modulation of the aging processes, Trumbull: Food and Nutrition Press, 1991: pp. 137-155.
- Weindruch R. Caloric restriction and aging. *Sci Am* 1996; 274: p.32-38.
- Anson RM. Absolute versus Relative Caloric Intake. Clues to the Mechanism of Calorie/Aging-Rate Interactions. *Ann N Y Acad Sci* 2004; 1019: 427-429.
- Ingram DK. Effects of dietary restriction on brain and behavioral function in aging rodents. In: Ingram DK, Baker GT, Shock NW, eds. The potential for nutrition modulation of the aging processes, Trumbull: Food and Nutrition Press, 1991: pp. 289-310.
- Socci DJ, Crandall BM, Arendash GW. Chronic antioxidant treatment improves the cognitive performance of aged rats. *Brain Res* 1995; 693: 88-94.
- Forster MJ, Dubey A, Dawson KM, et al. Age-related losses of cognitive function and motor skills in mice are associated with oxidative protein damage in the brain. *Proc Soc Exp Biol Med* 1996; 93: 4765-4769.
- Wu A, Wan F, Sun X, et al. Effects of dietary restriction on growth, neurobehavior, and reproduction in developing Kunming mice. *Toxicol Sci* 2002; 70: 238-244.
- Wu A, Sun X, Liu Y. Effects of caloric restriction on cognition and behavior in developing mice. *Neurosci Lett* 2003; 339: 166-168.
- Maswood N, Young J, Tilmont E, et al. Caloric restriction increases neurotrophic factor levels and attenuates neurochemical and behavioral deficits in a primate model of Parkinson's disease. *PNAS* 2004; 101: 18171-18176.
- De Oliveira LM. Malnutrition and environment: interaction effects upon animal behavior. *Revista Chilena de Nutrición* 1985; 13: 99-108.
- Almeida SS, Garcia RA, De Oliveira LM. Effects of early protein: malnutrition and repeated testing upon locomotor and exploratory behaviors in the elevated plus-maze. *Physiol Behav* 1993; 34: 749-752.
- Almeida SS, De Oliveira LM. Acquisition and extinction of jumping, two-way shuttle-box and bar press avoidance responses in malnourished rats: effects of shock intensity. *Braz J Med Biol Res* 1994; 27: 2443-2452.
- Santucci LB, Daud MM, Almeida SS, et al. Effects of early protein malnutrition and environmental stimulation upon the reactivity to diazepam in two animal models of anxiety. *Pharmacol Biochem Behav* 1994; 49: 393-398.
- Singh U, Devaraj S. Vitamin E: Inflammation and atherosclerosis. *Vitamins and Hormones* 2007; 76: 519-549.
- Hathcock JN, Azzu A, Blumberg J, Bray T, et al. Vitamins E and C are safe across a broad range of intakes. *Am J Clin Nutr* 2005; 81:736-745.
- Kontush A, Schekatolina S. Vitamin E in neurodegenerative disorders. Alzheimer's disease. *Ann N Y Acad Sci* 2004; 1031: 249-262.
- McDonald SR, Sohal RS, Forster MJ. Concurrent administration of coenzyme Q(10) and alpha-tocopherol improves learning in aged mice. *Free Radic Biol Med* 2005; 38: 729-736.
- Richwine AF, Godbout JP, Berg BM, et al. Improved psychomotor performance in aged mice fed diet high in antioxidants is associated with reduced ex vivo brain interleukin-6 production. *Brain Behav Immun* 2005; 19: 512-520.
- Oliveira SL, Diniz DB, Amaya-Farfan J. Carbohydrate energy restriction may protect the rat brain against oxidative damage and improve physical performance. *Brit J Nutr (London)* 2003; 89: 89-96.

20. Reeves PG, Nielsen FH, Fahey GC Jr. AIN-93 Purified diets for laboratory rodents: Final report of the American Institute of Nutrition Ad Hoc writing Committee on the reformulation of the AIN-76A Rodent Diet. *J Nutr* 1993; 123: 1939-1951.
21. Sharma A, Kumar A. Concurrent analysis of plasma retinol and α -tocopherol by isocratic HPLC. *Indian J Exp Biol* 1990; 28: 780-782.
22. Fernandez MS, Gonzalez C, Patterson AM. Oil enriched diets and behavioral parameters in rats' recovery from early under-nutrition. *Physiol Behav* 1997; 62: 113-119.
23. Moreira GMS, De Araújo M, De Oliveira LM, et al. The behavior of protein-calorie-malnourished rats on the elevated plus-maze test: an ethopharmacological analysis. *Psychobiology* 1997; 25: 180-185.
24. Riul TR, Almeida PS, Carvalho AF, et al. Effects of different levels of protein and environmental stimulation on the behavior of young rats tested in the elevated plus-maze. *Nutr Neurosci* 1998; 1: 295-303.
25. Almeida SS, Araújo M, Moreira GMS, et al. Short-term social isolation does not reduce elevated plus-maze exploration in early protein malnourished rats. *Nutr Neurosci* 1998; 1: 103-110.
26. Cadet JL, Katz M, Jackson-Lewis V, et al. Vitamin E attenuates the toxic effects of intrastriatal injection of 6-hydroxydopamine (6-OHDA) in rats: behavioral and biochemical evidence. *Brain Res* 1989; 476: 10-15.
27. Hass BS, Lewis SM, Duffy PH, et al. Dietary restriction in humans: report on the little rock conference on the value, feasibility, and parameters of a proposed study, *Mech Ageing Dev* 1996; 91: 79-94.
28. Weed JL, Lane MA, Roth GS, et al. Activity measures in rhesus monkeys on long-term calorie restriction. *Physiol Behav* 1997; 62: 97-103.
29. Joseph JA, Shukitt-hale B, Denisov NA, et al. Long-term dietary strawberry, spinach, or vitamin E supplementation retards the onset of age-related neuronal signal-transduction and cognitive behavioral deficits. *J Neurosci* 1998; 18: 8047-8055.
30. Shukitt-Hale B, Smith DE, Meydani M, et al. The effects of dietary antioxidants on psychomotor performance in aged mice. *Exp Gerontol* 1999; 34: 797-808.
31. Huston J, Burës J. Innate and motivated behavior. In: Burës J, Huston J, Buresöva O. *Techniques and basic experiments for the study of brain and behavior*, Netherlands: Elsevier, 1976: pp.49-59.
32. Sumien N, Heinrich KR, Sohal RS, et al. Short-term vitamin E intake fails to improve cognitive or psychomotor performance of aged mice. *Free Radic Biol Med* 2004; 36: 1424-1433.
33. Kim HK, Kim JH, Gao X, et al. Analgesic effect of vitamin E is mediated by reducing central sensitization in neuropathic pain. *Pain* 2006; 53-62.
34. Rodgers RJ, Cole JC. Anxiety enhancement in the murine elevated plus-maze by immediate prior exposure to social stress. *Physiol Behav* 1993; 53: 383-388.
35. Almeida SS, Soares EG, Bichuette MZ, et al. Effects of early postnatal malnutrition and chlordiazepoxide on experimental aversive situations. *Physiol Behav* 1992; 51: 1195-1199.
36. Goodrick CL, Ingram DK, Reynolds MA, et al. Differential effects of intermittent feeding and voluntary exercise on body weight and life span in adult rats. *J Gerontol* 1983; 38: 36-45.
37. Kryzhanovskii GN, Bakuleva LP, Luzina NL, et al. Role of the endogenous opioid system in the analgesic effect of α -tocopherol in dysmenorrhea. *Bull Exp Biol Med* 1988; 105: 162-164.
38. Kryzhanovskii GN, Luzina NL, Yarygin KN. α -tocopherol-induced activation of the endogenous opioid system. *Bull Exp Biol Med* 1989; 108: 1578-1579.

Recibido: 01-09-2008

Aceptado: 23-05-2009